

Probing the Character of the Pygmy Dipole Resonance

V. Derya^{1*}, J. Endres¹, M. N. Harakeh², D. Savran^{3,4},
H. J. Wörtche², and A. Zilges¹

¹*Institute for Nuclear Physics, University of Cologne, Germany*

²*KVI, Rijksuniversiteit Groningen, The Netherlands*

³*ExtreMe Matter Institute EMMI and Research Division, GSI, Darmstadt, Germany*

⁴*Frankfurt Institute for Advanced Studies FIAS, Frankfurt a.M., Germany*

Nuclear Structure 2012

**Argonne National Laboratory,
August 13 – 17, 2012**

Supported by the DFG (ZI 510/4-1 and ZI 510/4-2), by the EU under EURONS Contract No. RII3-CT-2004-506065 in the 6th framework programme, and by the Alliance Program of the Helmholtz Association (HA216/EMMI)




*Member of the Bonn-Cologne Graduate School of Physics and Astronomy


- Introduction
- Experimental Methods
- Results
 - for ^{94}Mo
 - for ^{48}Ca
- Summary

Introduction


Electric dipole strength in spherical atomic nuclei



Two-phonon
state




Electric pygmy
dipole resonance
(PDR)




Isovectorial electric
giant dipole resonance
(GDR)

Summed B(E1) Strength



Splitting of the PDR




- Real photons:
Strongly fragmented E1 strength

A. Zilges et al., Phys. Lett. B **542** (2002) 43
S. Volz et al., Nucl. Phys. **A779** (2006) 1
D. Savran et al., Phys. Rev. Lett **100** (2008) 232501

- Complementary α particles:
Splitting of the PDR
in ^{140}Ce , ^{138}Ba , and ^{124}Sn

J. Endres, E. Litvinova, *et al.*,
Phys. Rev. Lett. **105** (2010) 212503



Two Nuclei – Two Aspects

- ^{94}Mo


- Non-magic
- Near to (sub) shell closure
- Spherical

Effect of non-magicity?

- ^{48}Ca

- Doubly-magic
- Medium-mass region

Evolution of collectivity?



Experimental Methods

Reaction

- (γ, γ')
- Real-photon scattering

Setup

- Darmstadt High-Intensity Photon Setup
- $E_\gamma = 0\text{-}10\text{ MeV}$
- 2 HPGe detectors

Selectivity


- Mainly E1 from ground state

- $(\alpha, \alpha'\gamma)$
- Inelastic α -scattering with γ coincidence

- Big-Bite Spectrometer at KVI Groningen
- $E_\alpha = 136\text{ MeV}$
- 6-7 HPGe detectors and α spectrometer





- Mainly low spin from ground state
- Isoscalar probe

α - γ Coincidence Matrix




- Energy spectra through projection
- Selecting transitions by setting gates


Selecting transitions – Projected γ spectra




Results for ^{94}Mo



α -scattering
cross sections




$(\alpha, \alpha'\gamma)$ and (γ, γ') in ^{94}Mo



C. Romig,
private
communication

$(\alpha, \alpha'\gamma)$ and (γ, γ') in ^{140}Ce , ^{138}Ba , and ^{124}Sn



D. Savran *et al.*,
Phys. Rev. Lett. **97** (2006) 172502



J. Endres *et al.*,
Phys. Rev. C **80** (2009) 034302

J. Endres, E. Litvinova, *et al.*,
Phys. Rev. Lett. **105** (2010) 212503


- Splitting of the PDR \rightarrow due to isospin character
 - low-energy part $\rightarrow (\gamma, \gamma')$ and $(\alpha, \alpha'\gamma)$
 - high-energy part $\rightarrow (\gamma, \gamma')$ only

Low-Lying Dipole Strength in Lighter Nuclei

- Light-mass nuclei: halo nuclei, single-particle character excitations
- Medium-mass nuclei: development of a more collective dipole-excitation mode?
- Dependence on N/Z ratio in the Calcium chain





J=1⁻ states in ($\alpha, \alpha'\gamma$) and (γ, γ')



- Strongest state in (γ, γ') at 7.3 MeV is missing in ($\alpha, \alpha'\gamma$)
- Strongest state in ($\alpha, \alpha'\gamma$) at 7.6 MeV is weak in (γ, γ')

The State(s) at 7.655 MeV

- Known $J^\pi=3^-$ state at 7.651(1) MeV
- Multipolarity of the observed ground-state transition: E1 or E3?
- Double-differential cross section $d^2\sigma/(d\Omega_\alpha d\Omega_\gamma)$ and α - γ angular correlation \rightarrow Multipolarity





θ_γ : angle of γ -ray emission with respect to the α -beam (position of the 6 HPGe detectors)

Ground state transitions stem from $J^\pi=1^-$ state


Parity Assignments in ^{48}Ca

- Parity Measurement at the HI γ S facility
- 7 dipole excitations were observed



➔ The excited dipole states have **negative** parity

$J=1^-$ states in $(\alpha, \alpha'\gamma)$ and (γ, γ')



Summary

- Extension of the systematic study in (γ, γ') and $(\alpha, \alpha' \gamma)$ experiments by the nucleus ^{94}Mo
- Determination of branching ratios possible
- Dipole excitations in ^{48}Ca have been measured with three different experimental methods
- Strong octupole contribution to the strongest dipole excitation by α particles excluded
- Parity of the state excited by photons but not excited by α particles could be determined as negative



University of Cologne (Cologne, Germany)

J. Endres, A. Hennig, J. Mayer, L. Netterdon, S. Pascu,
S. G. Pickstone, A. Sauerwein, F. Schlüter, P. Scholz,
M. Spieker, T. M. Streit, and A. Zilges



Kernfysisch Versneller Instituut (Groningen, The Netherlands)

M. N. Harakeh and H. J. Wörtche



ExtreMe Matter Institute (Darmstadt, Germany)

D. Savran

Supported by:

